

# How does soil carbon sequestration affect greenhouse gas emissions from a sheep farming system? Results of a life cycle assessment case study

Pasquale Arca,<sup>1</sup> Enrico Vagnoni,<sup>1</sup> Pierpaolo Duce,<sup>1</sup> Antonello Franca<sup>2</sup>

<sup>1</sup>CNR-IBE, Institute of BioEconomy, National Research Council of Italy, Sassari; <sup>2</sup>CNR-ISPAAAM, Institute for Animal Production System in Mediterranean Environment, National Research Council of Italy, Sassari, Italy

## Highlights

- Extensification of dairy sheep systems provides an environmental benefit when soil C sequestration is considered.
- Extensification of dairy sheep systems determines lower environmental impact per hectare of utilized agricultural area.
- Enteric methane emissions are the main source of GHG emissions of the sheep milk life cycle.
- Carbon sequestration in permanent grasslands can considerably contribute to climate change mitigation.

## Abstract

A life cycle assessment (LCA) study of a transition from semi-intensive to semi-extensive Mediterranean dairy sheep farm suggests that the latter has a strong potential for offsetting greenhouse gas (GHG) emissions through the soil C sequestration ( $C_{seq}$ ) in permanent grasslands. The extensification process shows clear environmental advantage when emission intensity is referred to the area-based functional unit (FU). Several LCA studies reported that extensive livestock systems have greater GHG emissions per mass of product than intensive one, due to their lower productivity. However, these studies did not account for soil  $C_{seq}$  of temporary and permanent grasslands, that have a strong potential to partly mitigate the GHG balance of ruminant production systems. Our LCA study was carried out considering the transition from a semi-

intensive (SI) towards a semi-extensive (SE) production system, adopted in a dairy sheep farm located in North-Western Sardinia (Italy). Impact scope included enteric methane emissions, feed production, on-farm energy use and transportation, infrastructures as well as the potential C sink from soil  $C_{seq}$  compared to emission intensity. In order to provide a more comprehensive analysis, we used the following FUs: 1 kg of fat and protein corrected milk (FPCM) and 1 ha of utilised agricultural area (UAA). We observed that the extensification of production system determined contrasting environmental effects when using different FUs accounting for soil  $C_{seq}$ . When soil  $C_{seq}$  in emission intensity estimate was included, we observed slightly lower values of GHG emissions per kg of FPCM in the SI production system (from 3.37 to 3.12 kg CO<sub>2</sub> equivalents – CO<sub>2</sub>-eq), whereas a greater variation we observed in the SE one (from 3.54 to 2.90 kg CO<sub>2</sub>-eq). Considering 1 ha of UAA as FU and including the soil  $C_{seq}$ , the emission intensity in SI moved from 6257 to 5793 kg CO<sub>2</sub>-eq, whereas values varied from 4020 to 3299 kg CO<sub>2</sub>-eq in SE. These results indicated that the emission intensity from semi-extensive Mediterranean dairy sheep farms can be considerably reduced through the soil  $C_{seq}$ , although its measurement is influenced by the models used in the estimation.

Correspondence: Enrico Vagnoni, CNR-IBE, Institute of BioEconomy, National Research Council of Italy, Trav. La Crucca 3, 07100 Sassari, Italy. E-mail: enrico.vagnoni@ibe.cnr.it

Key words: Sheep farming; life cycle assessment; greenhouse gas emissions; carbon sequestration; extensification; permanent grasslands; temporary grasslands.

Acknowledgements: this study was supported by the LIFE financial instrument of the European Union (project SheepToShip LIFE - Looking for an eco-sustainable sheep supply chain: environmental benefits and implications, LIFE15 CCM/IT/000123).

Received for publication: 21 December 2020.

Revision received: 2 April 2021.

Accepted for publication: 8 April 2021.

© Copyright: the Author(s), 2021

Licensee PAGEPress, Italy

Italian Journal of Agronomy 2021; 16:1789

doi:10.4081/ija.2021.1789

This article is distributed under the terms of the Creative Commons Attribution Noncommercial License (by-nc 4.0) which permits any non-commercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

## Introduction

Sheep and goats represent about 60% of the total world ruminant population (FAO, 2019) and milk production is expected to increase globally in the next years (Pulina *et al.*, 2018). Sheep milk produced in Europe represents approximately 29% of global sheep milk production, with dairy sheep farms mainly concentrated in Mediterranean and Black Sea Regions (Pulina *et al.*, 2018). One of the largest producers in these areas is Sardinia (Italy), with more than 250,000 Mg year<sup>-1</sup>, which represents about 25% of total EU-27 sheep milk production (Rural Development Programme of Sardinia, 2014-2020). The main sheep breed raised in Sardinia is by far the autochthonous Sarda breed (Gutiérrez-Pena *et al.*, 2018), a dual-purpose breed (milk and meat) that can be considered one of the most important dairy sheep breeds in the world (Macciotta *et al.*, 1999) with more than 4.7 million heads reared in several Mediterranean areas. As well as in other Southern Europe areas, the Sardinian milk sheep sector is characterized by semi-

extensive farms where grazing on temporary and permanent grasslands is the main feeding source (Pulina *et al.*, 2018), with a wide range of natural resources and input utilization levels (Porqueddu *et al.*, 2017).

Sustainable intensification of production systems is clearly identified by scientists as key action for climate change mitigation strategy in agri-food sector (Gislon *et al.*, 2020). From this point of view, Sardinian sheep sector represents an interesting case study for testing strategies aimed to achieve a sustainable livestock supply chain and to better understand how to conciliate food provision with reduced environmental impacts. Several authors (Batalla *et al.*, 2015; Escribano *et al.*, 2020) showed how and to what extent the intensification level of Mediterranean dairy sheep farms affects the environmental performance of production systems, but the scientific evidences are not unambiguous and well defined. Usually, in life cycle assessment (LCA) studies, the intensification of agricultural and livestock systems improves the environmental performance per functional unit (FU) of product when the marginal yield increase is higher than the marginal input utilization (FAO, 2010; Notarnicola *et al.*, 2017), although the ecological optimum (eco-efficiency) depends on the specific situation (Hayashi *et al.*, 2006). On the other hand, in LCA studies on Mediterranean sheep and goat systems with different intensification levels, contrasting results are showed when soil C sequestration (soil  $C_{seq}$ ) is included in the emission intensity estimate per kg of normalized milk. Gutiérrez-Pena *et al.* (2019) and Batalla *et al.* (2015) showed that the emission intensity is not different in semi-intensive and semi-extensive systems when soil  $C_{seq}$  is included, whereas Escribano *et al.* (2020) observed higher environmental impact in semi-extensive ones, with or without soil  $C_{seq}$  inclusion. However, there is little agreement about the inclusion of the soil  $C_{seq}$  in LCA system boundaries. The methodological principle that excludes soil  $C_{seq}$  from LCA estimation is that C temporarily sequestered in the soil will be re-emitted in the future (Nayak *et al.*, 2019). In the recent Product Environmental Footprint Category Rules for dairy products (EDA, 2018), change in soil C level is considered as change in C stock and, consequently, excluded from the impact category 'Climate Change'. Nevertheless, some researchers have highlighted the importance of accounting for the soil C change in LCA studies, because C removed from the atmosphere and stored in the soil temporarily reduces the cumulative radioactive forcing over that time frame, reducing the climate impact (Levasseur *et al.*, 2013). Regarding these contrasting issues, Nayak *et al.* (2019) highlighted the availability of some estimation methods to be adapted to the context, but also confirmed what other authors pointed out about the unavailability of a common standard procedure for soil  $C_{seq}$  accounting in agricultural LCA (Brandão and i Canals, 2013; Petersen *et al.*, 2013; Arzoumanidis *et al.*, 2014).

Another scientific controversy within the agri-food LCA community concerns the criteria to be used to identify the most appropriate FU to express the environmental impacts of livestock systems. Salou *et al.* (2017) highlighted that the effects of intensification on emission intensity differ significantly depending on the FU adopted; indeed, the authors observed that the intensification does not produce variations on emission intensity per kg of normalized milk (mass FU), whereas higher values of emission intensity per ha (area-based FU) were observed in more intensive systems. In addition, Baldini *et al.* (2017) showed that the choice of the FU produces different results in terms of environmental output, advantaging in some case the more intensive systems and in other case the more extensive ones. Moreover, Escribano *et al.* (2020) stated that the use of the area-based FU and the inclusion of soil  $C_{seq}$  in the emission intensity estimate are more appropriate for environmental

assessment of extensive farming systems based on permanent grasslands. Finally, as observed by Gislon *et al.* (2020), few numbers of LCA studies on milk production are based on specific farm data and considered soil  $C_{seq}$  in the environmental profile of farming system. The main aim of this work was to evaluate the environmental implication of a Mediterranean dairy sheep farm, before and after the transition from a semi-intensive to a semi-extensive production system. For this purpose, we carried out an LCA study of a Sardinian dairy sheep farm including the contribution of soil  $C_{seq}$  and using both mass and area-based FUs.

## Materials and methods

In our research, a single case study approach was adopted. This approach is in accordance with the recommendations from Horrillo *et al.* (2021), who stated that the case study is a tool that allows to analyse in detail a specific phenomenon that occurred in a well-defined real context. Moreover, Fedele *et al.* (2014) demonstrated that an impacts assessment based on the LCA methodological approach can support a comparative environmental impact evaluation between contrasting production systems in a single farm. Ultimately, this methodology was suitable for our scope and, although it was limited to providing non-statistical results, it can contribute to scientific development with valid results when extrapolated from appropriate case studies (Flyvbjerg, 2006).

## Case study

The study was carried out in a dairy sheep farm located in Osilo, Sardinia (Italy) (40°45'11' N and 8°38'43' E, elevation 364 m a.s.l.) (Figure 1). The area has typical Mediterranean climate conditions with warm and dry summers, mild and wet winters (Chessa and Delitala, 1997). The average annual rainfall is approximately 760 mm, with 72 rainy days per year, mostly concentrated in October-November, and a drought period usually lasting from May to October. Minimum and maximum mean monthly temperatures are about 10 °C and 26 °C, respectively, with an average annual temperature of 16.5 °C. The landscape was characterized by hilly morphologies on volcanic rocks that occupied the fertile lowland, along a fluviokarstic valley that cuts a carbonate plateau (Biddau and Cidu, 2005). The surrounding area was characterized by dairy sheep farms with feed resources on tilled lands and grazing on permanent grasslands in areas unsuitable to crop production, which were neither inorganically fertilized nor irrigated, with small patches of native Mediterranean maquis. Permanent grasslands were grazed from autumn to spring, while temporary grasslands, such as annual crops, were grazed until the end of winter to allow the hay or grain production. Summer grazing was also carried out mainly on upland fields and on dry residuals of cereals and annual hay crops, after harvest/haymaking. During browsing, manure remained in the fields and contributed to replenish soil fertility levels. Native grasslands on neutral-subalkaline soils were characterized by annual species, with a dominant contribution of grasses.

Up until 2008, the case study farm was characterized by a foraging system based mainly on temporary grasslands such as grain-cereal crops (winter wheat, *Triticum durum* Desf.), annual forage crops (oat, *Avena sativa.*, and Italian ryegrass, *Lolium multiflorum* Lam.), both for grazing and hay production, and irrigated crop (silage maize, *Zea mays* L.) (Figure 1 and Table 1). The whole milk production was sold to the dairy industry. Since 2008, the farmer decided to change progressively the management strategy, with the

aim of destining all milk production to the on-farm manufacturing of semi-artisanal cheeses. As a result, most of the arable land has been converted from temporary grasslands to permanent grasslands, both natural and semi-natural (Figure 1 and Table 1). Natural grasslands were established exploiting the germination of native seedbank, while semi-natural grasslands through the over-seeding of annual self-reseeding legumes and grasses. The extensification of the production system, completed in 2011, was part of a wider farm management strategy oriented to increase the milk's added value, represented by the cheese selling, and to reduce its production costs (especially for self-produced forage). Therefore, this farm was characterized by the switching between two contrasting production systems that can be defined as 'semi-intensive' (SI, pre-2008) and 'semi-extensive' (SE, post-2008), respectively. The general characteristics of these dairy systems adopted by the farm during the two periods are reported in Table 2. Both farming systems used Sarda sheep breed. During the transition from SI to SE, the farmer slightly reduced flock size (340 and 320 productive ewes in SI and SE, respectively) and varied the animal diet (Tables 2 and 3). In the process of extensification, land management was changed (Table 1). The SI farming system managed 69.4 ha of utilized agricultural area (UAA), as temporary grasslands (30%, 35%, 22% and 9% for grazing, hay, grain and silage production, respec-

tively), and 3 ha as permanent grasslands (natural grassland for grazing) (Table 1). In the SE production system (69.7 ha), only 13% of the UAA was occupied by temporary grasslands (7% of Italian ryegrass-oat mixture and 6% of irrigated meadows - alfalfa, *Medicago sativa* L., and white clover, *Trifolium repens* L. - for grazing), while 87% was occupied by permanent grasslands, both natural and semi-natural (76% and 11%, respectively) (Table 1). Approximately 23% of natural grasslands were used for hay production, the remaining 77% for grazing. Semi-natural grasslands were used as grazing lands (Table 1). On-farm feed resources were integrated with about 92 and 85 Mg of concentrates in SI and SE, respectively.

### Life cycle assessment methodology

A comparative LCA was performed, according to the International Organization of Standardization LCA rules (14040 and 14044) (ISO, 2006a, 2006b). In order to have a more comprehensive view of the environmental impacts of sheep farming systems, soil  $C_{seq}$  was included in greenhouse gas (GHG) emissions balance, adopting two FUs: i) a mass-based FU, fat protein corrected milk (FPCM), where  $FPCM (kg) = raw\ milk (kg) \times (0.25 + 0.085\ fat\% + 0.035\ protein\%)$  (Pulina and Nudda, 2002); ii) an area-based FU, expressed in ha of UAA. The use of both FUs




Figure 1. Overview of the case-study farmland in North Sardinia (Italy). The two farming systems (semi-intensive, SI, and semi-extensive, SE) are overlaid with a land use map during the analysed period (based on satellite imageries of SardegnaFotoAeree: 2006 for semi-intensive system and 2013 for semi-extensive system).

allowed to combine productive and economic results with depletion of natural resources, reflecting, in other terms, the two main functions of agricultural production systems: the production of market goods and the provision of public services and externalities, associated with the environmental role of farming systems (Basset-Mens and van der Werf, 2005; Gutiérrez-Peña *et al.*, 2019). By this way, global and local effects of climate change were included in the perspective of the analysis, giving back a more balanced assessment of the results. Specific farm data were referred to the years 2001 and 2011, when the farm was characterized by two different production systems. The system boundary of the analysis was from 'cradle-to-farm-gate'. In particular, system boundary of the study included: i) amount of hay, green forage and concentrat-

ed consumed by the flock, comparing the biomass yields of grasslands and the nutritional needs of each animal category (based on gender, age, weight, physiological stage and production level of animals); ii) water and energy use; iii) machineries and equipment (tractors included); iv) milking parlour, barns and other manufactured goods linked with the farm structure; v) consumable materials (agrochemicals, packaging materials, *etc.*); vi) distances and mode of transportations. We collected primary data, representing more than 90% of the inventory data, through farm's register examination, several field visits and interviews to farmer. All representative secondary data were taken from Ecoinvent Centre v3.6 database (Moreno Ruiz *et al.*, 2018), except for the dataset of sunflower meal and soybean feed, taken from Agri-footprint 4.0

**Table 1. Grassland type, use of the biomass and surface area (ha) in the semi-intensive (SI) and semi-extensive (SE) farming systems.**

| Temporary or permanent grassland | Type of grassland                       | Semi-intensive system (SI)<br>Species or mixture   | Use     | Surface (ha) |
|----------------------------------|---|--|---------|--------------|
| Temporary grasslands             | Cereal crop<br>Annual forage crops      | Winter wheat   | Grain   | 16.0         |
|                                  |   | Oat  | Grazing | 2.0          |
|                                  |   | Italian ryegrass-oat mixture   | Grazing | 20.0         |
|                                  | Irrigated crop                          | Maize  | Hay     | 25.0         |
|                                  |   |  | Silage  | 6.4          |
| Permanent grassland              | Natural grassland                       | Native vegetation  | Grazing | 3.0          |
| Temporary or permanent grassland | Type of grassland                       | Semi-extensive system (SE)<br>Species or mixture   | Use     | Surface (ha) |
| Temporary grasslands             | Annual forage crop<br>Irrigated meadows | Italian ryegrass-oat mixture   | Grazing | 4.9          |
|                                  |   | Alfalfa  | Grazing | 2.6          |
|                                  |   | White clover   | Grazing | 1.7          |
| Permanent grasslands             | Semi-natural grasslands                 | Type I: mixture of <i>Lolium rigidum</i> ,<br><i>Trifolium subterraneum</i>                                | Grazing | 1.0          |
|                                  |   | Type II: mixture of <i>Lolium rigidum</i> ,<br><i>Medicago polymorpha</i>                                  | Grazing | 1.9          |
|                                  |   | Type III: mixture of <i>Lolium rigidum</i> , <i>Medicago polymorpha</i> ,<br><i>Trifolium subterraneum</i> | Grazing | 4.9          |
|                                  | Natural grassland                       | Native vegetation  | Grazing | 40.6         |
|                                  |   |  | Hay     | 12.1         |

**Table 2. Main characteristics of the two production systems, namely semi-intensive (SI) and semi-extensive (SE), adopted on the same farm in different years.**

| Characteristics of dairy sheep farm                                  | UM   | SI      | SE     |
|--|--|---------|--------|
| Heads (number of mature ewes)*                                       | n  | 340     | 320    |
| Stocking rate*   | head ha <sup>-1</sup>                      | 4.7     | 4.6    |
| Milk total annual production   | kg   | 104,234 | 82,214 |
| Milk per capita annual production                                    | kg ewe <sup>-1</sup> year <sup>-1</sup>    | 307     | 257    |
| Fat and protein corrected milk (FPCM), per capita annual production  | kg ewe <sup>-1</sup> year <sup>-1</sup>    | 303     | 227    |
| Dry matter intake (DMI) of mature ewes*                              | kg DM ewe <sup>-1</sup> year <sup>-1</sup> | 653     | 565    |
| Fraction of concentrate in the diet of mature ewes*                  | kg DM DMI <sup>-1</sup>                    | 0.28    | 0.31   |
| Temporary grasslands (cereal and annual forage crops)                | ha   | 63.0    | 4.9    |
| Temporary grasslands (irrigated silage maize)                        | ha   | 6.4     | 0      |
| Temporary grasslands (irrigated meadows of alfalfa and white clover) | ha   | 0       | 4.3    |
| Permanent grasslands (natural and semi-natural)                      | ha   | 3.0     | 60.5   |
| Mineral N-fertilizing  | kg ha <sup>-1</sup>                        | 68.9    | 0.5    |
| Mineral P <sub>2</sub> O <sub>5</sub> -fertilizing                   | kg ha <sup>-1</sup>                        | 104.4   | 1.7    |

\*Rams and replacement are not considered.

(2017) database. No generic data were used. The estimation of enteric CH<sub>4</sub> emission (F<sub>CH<sub>4</sub></sub>) was based on the use of the CH<sub>4</sub> emission factor (Y<sub>m</sub>), calculated as function of Metabolizable Energy Intake (MEI) (Vermorel *et al.*, 2008):

$$F_{CH_4} = MEI \times Y_m / 55.65 \quad (1)$$

where:

F<sub>CH<sub>4</sub></sub> represented the kg of emitted CH<sub>4</sub>/day per head; MEI was expressed in MJ/day per head; the coefficient 55.65 represented the energy content of 1 kg of CH<sub>4</sub> and was expressed in MJ; Y<sub>m</sub>, the methane conversion factor (%), which expressed the proportion of ration gross energy lost as CH<sub>4</sub>, was calculated as:

$$Y_m = -0.15 \times DE + 21.89 \quad (2)$$

where DE was the Digestible Energy (DE in %) of the diet.

The monthly diets of each animal category were defined through farm data collection. Type and amount of feed utilized in the diet were collected as primary data, except for the intake of grazed biomass. For the composition of each feed type we used the database elaborated by LAORE Sardegna (the Regional Agency for Agriculture Development) (Sardegna Agricoltura, 2013). In order to calibrate the diet, based on the nutritional needs of each animal category and the nutritional value of each feed type, the Small Ruminant Nutrition System Software (SRNS - Tedeschi *et al.*, 2008) was adopted using amount, type and composition of the

**Table 3. Main input (feed, diesel, electricity and water consumptions) and output (feed production and feed self-sufficiency) in semi-intensive (SI) and semi-extensive (SE) production systems, adopted on the same farm in different years.**

| Item                                      | Unit                                       | Value  |        |
|---|--|--------|--------|
|   |  | SI     | SE     |
| <b>Input</b>                              |  |        |        |
| Hay/silage consumption*                   | kg DM ewe <sup>-1</sup> year <sup>-1</sup> | 314    | 70     |
| Green forage consumption*                 | kg DM ewe <sup>-1</sup> year <sup>-1</sup> | 155    | 320    |
| Concentrate consumption*                  | kg DM ewe <sup>-1</sup> year <sup>-1</sup> | 184    | 176    |
| Purchased feed per kg FPCM                | kg DM kg FPCM <sup>-1</sup>                | 0.90   | 1.18   |
| Unit diesel consumption <sup>o</sup>      | kg diesel 100 kg FPCM <sup>-1</sup>        | 6.93   | 4.56   |
| Unit electricity consumption <sup>#</sup> | kWh 100 kg FPCM <sup>-1</sup>              | 0      | 9.75   |
|   | kg diesel 100 kg FPCM <sup>-1</sup>        | 6.71   | 0      |
| Unit water consumption                    | m <sup>3</sup> 100 kg FPCM <sup>-1</sup>   | 19.67  | 11.48  |
| <b>Output</b>                             |  |        |        |
| Hay/silage production                     | Mg DM year <sup>-1</sup>                   | 120.49 | 26.48  |
| Grazing production                        | Mg DM year <sup>-1</sup>                   | 60.77  | 110.13 |
| Concentrate production                    | Mg DM year <sup>-1</sup>                   | 22.5   | 0      |
| Feed self-sufficiency <sup>s</sup>        | %  | 69     | 62     |

\*Consumptions of rams and replacement are not included; <sup>o</sup>diesel use includes the fossil fuel consumption for agricultural operations; <sup>#</sup>electricity use expressed as kg of fossil fuel from power generator in SI and kWh from electric company in SE; <sup>s</sup>ratio feed production/feed consumption.

**Table 4. Annual biomass yield and total residues (above- and belowground) for each grassland type in both production systems adopted on the same farm in different years.**

| Item                                   | Yield (Mg DM ha <sup>-1</sup> ) | Total residues (Mg DM ha <sup>-1</sup> ) |
|--|---------------------------------|--|
| Semi-intensive system                  |                                 |  |
| Oat (grazing)                          | 2.12                            | 2.55                                     |
| Italian ryegrass-oat mixture (grazing) | 2.41                            | 3.33                                     |
| Italian ryegrass-oat mixture (hay)     | 1.01                            | 1.23                                     |
| Maize (silage)                         | 14.61                           | 4.02                                     |
| Winter wheat (grain)                   | 1.41                            | 3.28                                     |
| Natural grassland (grazing)            | 2.77                            | 11.36                                    |
| Semi-extensive system                  |                                 |  |
| Italian ryegrass-oat mixture (grazing) | 5.57                            | 7.69                                     |
| Alfalfa (grazing)                      | 10.2                            | 15.52                                    |
| White clover (grazing)                 | 2.21                            | 11.94                                    |
| Semi-natural grassland (grazing)*      | 3.06                            | 8.12                                     |
| Natural grassland (grazing)            | 0.66                            | 2.72                                     |
| Natural grassland (hay)                | 2.18                            | 7.71                                     |

\*Mean values of the three type of semi-natural grasslands are reported.

feed as inputs. The intake of grazed biomass was estimated by difference between total dry matter intake (DMI), calculated with the SRNS software, and the sum of the amounts of the other feeds.

Emissions related to pesticide and fertilizer applications were assessed according to the following approaches: equations reported in Ecoinvent report No.15 (Nemecek and Kägi, 2007) for: i) emissions of  $\text{NO}_x$  to air; ii) emissions of heavy metals,  $\text{PO}_3^-$ , P and  $\text{NO}_3^-$  to water; and iii) emissions of heavy metals to soil; Tier 1 IPCC method (IPCC, 2019) for both  $\text{N}_2\text{O}$  direct and indirect and  $\text{CO}_2$  emissions to air; Tier 2 IPCC method (IPCC, 2019), using national emission factor proposed by ISPRA (2011) for  $\text{NH}_3$  emissions to air. The impacts related to manure management excluded  $\text{CH}_4$  emissions and included only the  $\text{N}_2\text{O}$  emitted through animal excreta, with the rationale that in both farming systems sheep were not confined in small or covered spaces. This type of animal emission was estimated following the IPCC (2019) approach and using the default emission factor for sheep and 'other animals' [ $0.003 \text{ kg N}_2\text{O-N (kg N)}^{-1}$ ]. In addition, daily N excretion of animal categories was estimated based on empirical equations (Decandia *et al.*, 2011) for ewes (lactating, dry, pregnant and replacements), rams and lambs. Final and intermediate transports were inventoried considering means of transport, distances and transported mass. To calculate distances, primary data were used when available (internet researches were done to find production plants and logistic chain). Big size machineries road transport was modelled referring to the corresponding Ecoinvent processes. In case of lack of primary data, logistic and distances were traced utilizing Searates website (Searates, 2021). Fossil fuel consumptions were estimated by adding up consumptions of all the agricultural operations. For the year 2001, fossil fuels consumption included also the use of the power generator. Electricity consumption for the year 2011 (in the year 2001 the farm used an electric generator as power supply) was calculated considering the average annual consumption reported in the supply bills of the electric company, excluding consumption for family and external uses. In addition, consumptions of major utilities (such as irrigation, milking and milk refrigeration) were estimated based on installed power and by checking literature data. Finally, the estimated data were compared with those reported in the bills to identify any discrepancies. Electricity datasets were built based on the energetic mix declared by the elec-

tric company for the reference year, starting from the Ecoinvent process 'Electricity, high voltage {IT} market for | Cut-off, U'.

In line with several LCA investigations on dairy sector (Pirlo *et al.*, 2014; Baldini *et al.*, 2017), we performed an economic allocation procedure in order to partitioning all inputs and outputs, considering that: i) milk, the 'main' product, had a very higher economic value than co-products such as meat, live rams (only in SI system) and wool; ii) in similar cases, allocation mode did not affect the LCA results (Salou *et al.*, 2017). In SI system, the economic allocation resulted as follows: 76% to milk, 13% to rams, 10% to meat and 1% to wool. Similarly, in SE system, it resulted in 91% to milk, 8% to meat and 1% to wool. We used SimaPro software (PRé Consultants, 2018) to model the life cycle and for impact analysis. The LCA analysis focused exclusively on Climate Change impact category, expressed as emission intensity. We calculated emission intensity using the IPCC (2013) evaluation method, based on Global Warming Potential (GWP) indicator (100-year time horizon) expressed in kg of  $\text{CO}_2$  equivalents ( $\text{CO}_2\text{-eq}$ ), and with the latest values of  $\text{CH}_4$  characterization factor (34.00 and  $36.75 \text{ kg CO}_2\text{-eq/kg}$  for biogenic and fossil  $\text{CH}_4$ , respectively).

### Estimation of soil carbon sequestration

We calculated the soil  $C_{\text{seq}}$ , referring to the total area actually used to feed sheep (both for grain and forage supply), according to the method suggested by Petersen *et al.* (2013). This model was designed exclusively for agricultural LCA studies for estimating the soil C changes as a consequence of the C input from above- and belowground crop residues and manure added to the soil. This approach was based on the modelling of two C fluxes: i) from the soil to the atmosphere, where the soil organic matter mineralization was modelled using the soil C model C-TOOL (Petersen, 2010); ii) from the atmosphere to the soil, where the atmospheric  $\text{CO}_2$  decay was modelled using the Bern Carbon Cycle model (IPCC, 2007). Petersen *et al.* (2013) observed that 9.7% of C added to the soil as organic C input in the first year would be sequestered in a 100-year perspective. This method, although being simplistic, was based on site-specific data of soil C input and field conditions, whereas the other available models to estimate the soil  $C_{\text{seq}}$  in agricultural LCA were based on default values per ha of grassland, as

**Table 5. Equations and coefficients used to estimate the aboveground residues depending on grassland type and use destination of the biomass.**

| Type of grassland  | Use     | Equation   | Acronym   | Index                              | Source                        |
|--|---------|--|---|------------------------------------|-------------------------------|
| Temporary grasslands:<br>- annual forage crops*<br>- irrigated meadows <sup>o</sup>    | Grazing | $\text{AbRes}_{\text{grz}} = \text{TAbB}_{\text{grz}} \times i_{\text{grz}}$   | <ul style="list-style-type: none"> <li>• <math>\text{AbRes}_{\text{grz}}</math> = aboveground residues after sheep grazing</li> <li>• <math>\text{TAbB}_{\text{grz}}</math> = total aboveground biomass of grazed grassland</li> <li>• <math>\text{Yic}_{\text{grz}}</math> = dry yield (green forage) of grazed grassland (available data)</li> <li>• <math>i_{\text{grz}}</math> = index of aboveground residues after sheep grazing</li> </ul> | $i_{\text{grz}} = 0.25$            | Seddaiu <i>et al.</i> (2018)  |
| Permanent grasslands:<br>- semi-natural grasslands <sup>#</sup><br>- natural grassland |         | $\text{TAbB}_{\text{grz}} = \text{Yic}_{\text{grz}} \times (1 - i_{\text{grz}})^{-1}$                                      |   |                                    |                               |
| Temporary grasslands:<br>- cereal crops <sup>§</sup>                                   | Grain   | $\text{AbRes}_{\text{grain}} = (\text{Yic}_{\text{grain}} \times \text{HI}_{\text{cer}}^{-1}) - \text{Yic}_{\text{grain}}$ | <ul style="list-style-type: none"> <li>• <math>\text{AbRes}_{\text{grain}}</math> = aboveground residues after grain harvest</li> <li>• <math>\text{Yic}_{\text{grain}}</math> = dry grain yield of cereal (available data)</li> <li>• <math>\text{HI}_{\text{cer}}</math> = harvest index of cereal (winter wheat)</li> </ul>  | $\text{HI}_{\text{cer}} = 0.4322$  | Bolinder <i>et al.</i> (1997) |
| Permanent grasslands:<br>- natural grassland   | Hay     | $\text{AbRes}_{\text{shay}} = \text{Yic}_{\text{hay}} \times i\text{LO}_{\text{shay}}$                                     | <ul style="list-style-type: none"> <li>• <math>\text{AbRes}_{\text{shay}}</math> = aboveground residues after hay harvest</li> <li>• <math>\text{Yic}_{\text{hay}}</math> = dry hay yield of natural pasture (available data)</li> <li>• <math>i\text{LO}_{\text{shay}}</math> = index of hay harvest losses</li> </ul>   | $i\text{LO}_{\text{shay}} = 0.185$ | Lai <i>et al.</i> (2017)      |

\*Oat and Italian ryegrass-oat mixture; <sup>o</sup>Alfalfa and white clover; <sup>#</sup>mixtures of annual self-seeding species (rigid ryegrass, burr medic and subterranean clover); <sup>§</sup>winter wheat.

highlighted by Batalla *et al.* (2015). In addition, the 100-year time horizon was in line with the time perspective of GWP indicator. Therefore, other authors used this method to estimate soil  $C_{seq}$  in LCA studies on dairy systems under Mediterranean conditions (Batalla *et al.*, 2015; Gutiérrez-Peña *et al.*, 2019; Escribano *et al.*, 2020), as well as in regions of Western Europe (Knudsen *et al.*, 2019). In order to estimate soil  $C_{seq}$ , the same coefficient (9.7%) was applied to the amount of soil C input, composed of two distinct fractions: i) the C derived from crop residues; and ii) the C contained in manure deposited by sheep during grazing (Batalla *et al.*, 2015). The C derived from crop residues included both C from aboveground crop residues and from belowground biomass, left on the soil at the end of the first year. The estimation of above- and belowground residues was based on the available data of each grassland yield, expressed in Mg of dry matter (DM)  $ha^{-1}$  (Table 4). For all grasslands, we converted the amounts of residues into C using a coefficient of C content equal to 0.40 (Burle *et al.*, 1997; dos Santos *et al.*, 2011), except for silage maize and Italian ryegrass-oat mixture destined to the hay production, for which we estimated the C derived from crop residues as percentage of the harvested DM, 11% and 44.7%, respectively, following Lai *et al.* (2017). Aboveground residues were estimated using different equations, by applying coefficients to the total aboveground biomass or yield, depending on the grassland type and use destination of the biomass, as reported in Table 5. The belowground residues included roots and rhizodeposition biomass. We computed the root biomass by applying a specific shoot-root or root-shoot ratio index (see Table 6 for literature details) to the relative total aboveground biomass, estimated for each grassland as sum of yield and aboveground residues. Rhizodeposition was calculated as fraction of the entire root system biomass, using an index equal to 0.65 (Bolinder *et al.*, 2007) for each grassland. In natural grasslands and irrigated meadows (alfalfa and white clover), we estimated roots as annual biomass increase and rhizodeposition as a fraction of the

entire root system. By contrast, in semi-natural grasslands and temporary grasslands, such as cereal crops and annual forage crops, both roots and rhizodeposition were estimated as annual biomass production. Finally, C:N ratio index equal to 13.4 (Escudero *et al.*, 2012) was used for estimating the amount of C input derived from sheep manure during grazing.

## Results

### Farming systems

Both SE and SI farming systems had almost the same number of lactating ewes (320 *versus* 340 heads) and slightly different values of stocking rate (4.6 *versus* 4.7 head  $ha^{-1}$ ) (Table 2). During transition to SE, the diet of productive ewes varied (Table 3), with an increase of the green forage fraction and a reduced use of conserved forages (0.57 and 0.12 kg  $DMI^{-1}$  in SE, 0.24 and 0.48 kg  $DMI^{-1}$  in SI, respectively). The DMI of ewes in SE was lower for about 88 kg DM  $ewe^{-1}$  (565 *versus* 653 kg DM  $ewe^{-1}$  year $^{-1}$  in SE and SI, respectively) (Table 2). The fraction of concentrates in the diet slightly increased from SI to SE (0.28 and 0.31 kg concentrate kg  $DMI^{-1}$ ) (Table 2), while individual concentrate consumptions of ewes were similar in both farming systems (184 *versus* 176 kg  $ewe^{-1}$  year $^{-1}$ ) (Table 3). Consequently, ewe milk productivity was lower in SE compared with SI (227 *versus* 303 kg FPCM  $ewe^{-1}$  year $^{-1}$ ) (Table 2). Feed efficiency (FE) of productive ewes was lower in SE than SI: 0.40 *versus* 0.46 kg FPCM kg  $DMI^{-1}$ , respectively. Annual feed production in SE was about 67% of SI (Figure 2), while annual FPCM production was lower by 30% (Table 2). Furthermore, despite the lower amount of total purchased feeds (85.33 *versus* 97.38 Mg DM year $^{-1}$  in SE and SI, respectively), feed self-sufficiency in SE was lower than in SI (62% *versus* 68%, respectively) (Table 3).

**Table 6. Equations and coefficients applied to estimate the belowground residues depending on the type and duration time of the grassland.**

| Type of grassland                       | Duration (year) | Equation   | Acronym  | Index   |
|---|-----------------|--|--|---|
| Temporary grasslands:                   |                 |  |  |   |
| i) annual forage crops                  |                 |  | <ul style="list-style-type: none"> <li>BelRes<sub>an</sub> = belowground residue of annual crop</li> <li>i(S:R)<sub>an</sub> = shoot-root ratio index of annual crop</li> <li>iRz = index of Rhizodeposition as fraction of root biomass</li> <li>TAAbB = total aboveground biomass*</li> </ul>  | <ul style="list-style-type: none"> <li>i(S:R)<sub>an</sub> = 2.53 (oat)<sup>o</sup></li> <li>i(S:R)<sub>an</sub> = 3.76 (winter wheat)<sup>o</sup></li> <li>i(S:R)<sub>an</sub> = 2.1 (Italian ryegrass-oat mixture)<sup>#</sup></li> <li>iRz = 0.65<sup>o</sup></li> </ul> |
| - oat                                   | 1               | BelRes <sub>an</sub> = [TAAbB × i(S:R) <sub>an</sub> <sup>-1</sup> × (1+iRz)]                                      |  |   |
| - Italian ryegrass-oat mixture (grazed) | 1               |  |  |   |
| ii) cereal crop                         |                 |  |  |   |
| - winter wheat                          | 1               |  | <ul style="list-style-type: none"> <li>BelRes<sub>per</sub> = belowground residue of perennial crop</li> <li>i(R:S)<sub>per</sub> = root-shoot ratio index of perennial crop</li> <li>t = time of crop duration</li> <li>iRz = index of Rhizodeposition as fraction of root biomass</li> <li>TAAbB = total aboveground biomass*</li> </ul>     | <ul style="list-style-type: none"> <li>i(R:S)<sub>per</sub> = 0.9901 (alfalfa)<sup>o</sup></li> <li>i(R:S)<sub>per</sub> = 4.224 (white clover and natural pasture)<sup>§</sup></li> <li>iRz = 0.65<sup>o</sup></li> </ul>  |
| Temporary grasslands:                   |                 |  |  |   |
| i) irrigated meadows                    |                 |  |  |   |
| - alfalfa                               | 4               | BelRes <sub>per</sub> = [TAAbB × i(R:S) <sub>per</sub> × t <sup>-1</sup> ] + [TAAbB × i(R:S) <sub>per</sub> × iRz] |  |   |
| - white clover                          | 4               |  |  |   |
| Permanent grasslands:                   |                 |  |  |   |
| i) natural grassland                    | 50              |  | <ul style="list-style-type: none"> <li>BelRes<sub>ss</sub> = belowground residue of self-seeding crop</li> <li>i(R:S)<sub>ss</sub> = root-shoot ratio index of self-seeding crop</li> <li>t = time of crop duration</li> <li>iRz = index of Rhizodeposition as fraction of root biomass</li> <li>TAAbB = total aboveground biomass*</li> </ul> | <ul style="list-style-type: none"> <li>i(R:S)<sub>ss</sub> = 4.224<sup>§</sup></li> <li>iRz = 0.65<sup>o</sup></li> </ul>   |
| Permanent grasslands:                   |                 |  |  |   |
| i) semi-natural grassland               |                 |  |  |   |
| - annual self-seeding mixtures          | 4               | BelRes <sub>ss</sub> = [TAAbB × i(R:S) <sub>ss</sub> × t <sup>-1</sup> ] × (1 + iRz)                               |  |   |

\*TAAbB = AbRes + Yie, for each grassland Total Aboveground Biomass (TAAbB) is estimated summing Aboveground Residues (AbRes, Table 5) and Yield (Yie, Table 4); <sup>o</sup>Bolinder *et al.*, 2007; <sup>#</sup>Lai *et al.*, 2017; <sup>§</sup>Mokany *et al.*, 2005.

## Life cycle assessment

As reported in Table 7, by excluding soil  $C_{seq}$ , we observed higher values of emission intensity per kg FPCM in SE than in SI (for about 5%), whereas opposing results were obtained using the area-based FU. Specifically, the emission intensity per kg FPCM was equal to 3.54 and 3.37 kg  $CO_2$ -eq kg FPCM<sup>-1</sup> in SE and SI, respectively. However, considering the area-based FU, the emission intensity assessed without soil  $C_{seq}$  was lower in SE than in SI (4030 *versus* 6257 kg  $CO_2$ -eq ha UAA<sup>-1</sup>, respectively).

In the calculation of emission intensity that includes the contribution of soil  $C_{seq}$ , the GHG emissions per kg of FPCM and per ha of UAA showed a reduction by about 18% and 7% in SE and SI farming systems, respectively (Table 7). The values of emission intensity including soil  $C_{seq}$  were equal to 2.90 *versus* 3.12 kg  $CO_2$ -eq kg FPCM<sup>-1</sup> and 3299 *versus* 5793 kg  $CO_2$ -eq ha UAA<sup>-1</sup> in SE and SI, respectively.

The contribution analysis (Table 7) showed animal emissions as the main source of GHG emissions, with much more than 50% of contribution in both farming systems. Animal emissions included enteric  $CH_4$  and faecal  $N_2O$  emissions, and the latter were by far the lowest impacting, representing less than 1% of the total emission intensity. Immediately after animal emissions, off-farm and on-farm feed productions accounted for a relevant hotspot, representing on average more than 19% of the total emission intensity in both production systems, including and excluding soil  $C_{seq}$  (Table 7). As expected, the contribution of purchased feed was higher in the SE farm management and the difference between the two production systems mainly concerned the role of on-farm feeds, since in SI represented on average 15% of the total feed contribution, while in SE accounted by only 1%.


## Carbon sequestration in soil

In the transition from SI to SE dairy sheep system, we estimated changes in soil  $C_{seq}$ , with a value in SE almost twice than in SI (55.30 *versus* 33.90 Mg  $CO_2$  year<sup>-1</sup>), although the total grasslands area actually utilized in both systems was similar (69.7 ha *versus* 72.4 ha) (Figures 2 and 3). Annual soil  $C_{seq}$  from manure during grazing was similar in both farming systems (2.57 *versus* 3.36 Mg  $CO_2$  year<sup>-1</sup> in SE and SI, respectively). However, annual soil  $C_{seq}$  from crop residues was higher in SE than SI (52.73 *versus* 30.55

Mg  $CO_2$  year<sup>-1</sup>, respectively), representing the main input of the total soil  $C_{seq}$  (Figure 2).

Taking into account the grassland use destination, the soil  $C_{seq}$  contribution of grazed grasslands in SI was almost the same of grasslands for hay, silage and grain purposes (15.06 *versus* 15.49 Mg  $CO_2$  year<sup>-1</sup>), while in SE soil  $C_{seq}$  from grazed grasslands was 39.38 Mg  $CO_2$  year<sup>-1</sup>, accounting for 75% of the total (Figure 3).

In SI, natural permanent grasslands contributed for about 16% of soil  $C_{seq}$  from crop residues, even though they covered only 4% of UAA (Figure 3). By contrast, in SE the amount of soil  $C_{seq}$  due to permanent grasslands (both natural and semi-natural) and irrigated meadows (alfalfa and white clover) contributed for about 90% of the soil  $C_{seq}$  from crop residues, while temporary grasslands such as annual forage crops sequestered the remaining 10% (Figure 3).



**Figure 2.** Feed production and soil C sequestration. Total on-farm feed production (Mg DM of grain and forages) and soil C sequestration (Mg  $CO_2$ ) from crop residues and manure in the semi-intensive and semi-extensive systems. The label values indicate the total on farm feed production (Mg DM).

**Table 7.** Process contributions for the Climate Change impact category. Emission and contribution of the processes to the total greenhouse gas emissions of semi-intensive (SI) and semi-extensive (SE) production systems, calculated including and excluding soil C sequestration (soil  $C_{seq}$ ), for both 1 kg of fat and protein corrected milk (FPCM) and 1 ha of utilised agricultural area (UAA) functional units.

| Climate Change                                     | Soil $C_{seq}$ excluded |      | Soil $C_{seq}$ included |      |
|--|-------------------------|------|-------------------------|------|
|  | SI                      | SE   | SI                      | SE   |
| kg $CO_2$ -eq per kg FPCM                          | 3.37                    | 3.54 | 3.12                    | 2.90 |
| kg $CO_2$ -eq per ha UAA                           | 6257                    | 4030 | 5793                    | 3299 |
| Process contribution (%)                           |                         |      |                         |      |
| Animal emissions                                   | 56                      | 65   | 61                      | 80   |
| Purchased feeds                                    | 12                      | 18   | 13                      | 22   |
| On-farm feeds                                      | 15                      | 1    | 16                      | 1    |
| Power supply                                       | 6                       | 3    | 7                       | 4    |
| Transport (lorry and/or transoceanic freight ship) | 3                       | 4    | 3                       | 5    |
| Infrastructures                                    | 1                       | 0    | 1                       | 0    |
| Tractor and agricultural machinery production      | 0                       | 3    | 0                       | 3    |
| Soil C sequestration                               | 0                       | 0    | -8                      | -22  |
| Remaining processes*                               | 7                       | 6    | 7                       | 7    |

\*All processes with a percentage contribution lower than 0.35% are included.

Overall, by referring soil  $C_{seq}$  to 1 kg of FCPM, the values were 0.76 and 0.33 kg of  $CO_2$  sequestered in SE and SI, respectively. When soil  $C_{seq}$  was referred to the unit of area, the values were equal to 793 and 469 kg  $CO_2$  per ha of UAA.

## Discussion

### Life cycle assessment


The extensification of the production system showed contrasting environmental effects, depending on the FU used and the inclusion or exclusion of soil  $C_{seq}$  in LCA system boundaries (Table 7). Gutiérrez-Peña *et al.* (2019) and Escribano *et al.* (2020) reported similar findings in a comparison between Spanish dairy goat and sheep systems with different grazing regimes, respectively.

Excluding soil  $C_{seq}$ , the difference of emission intensity per kg of FPCM between SI and SE can be explained by the decrease of FPCM production in SE. This observation is consistent with data from previous studies that showed how extensive livestock systems have a greater environmental impact than intensive systems, due to their less productive and less efficient management (Gerber *et al.*, 2013). Moreover, by conducting the LCA analysis using the kg FPCM<sup>-1</sup> as FU and excluding soil  $C_{seq}$ , our results are similar to those reported by other investigations (Atzori *et al.*, 2015; Vagnoni *et al.*, 2017) and comparable with those observed by Batalla *et al.* (2015). The latter, who carried out a study in dairy sheep systems under similar conditions to ours, found emission intensity ranging

from 2.87 to 3.19 kg  $CO_2$ -eq kg FPCM<sup>-1</sup> in three semi-intensive systems, and from 2.76 to 5.17 kg  $CO_2$ -eq kg FPCM<sup>-1</sup> in six semi-extensive systems. However, our results differ, in some aspects, from those of Escribano *et al.* (2020), who stated that emission intensity values followed a trend corresponding to the increasing level of extensification (four farming systems in total), ranging from 1.77 (most intensive farm) to 4.09 (most extensive farm) kg  $CO_2$ -eq kg FPCM<sup>-1</sup>. This lower emission intensity per kg of FPCM could be associated to the lower values of enteric  $CH_4$  emissions factors (25 instead of 34 kg  $CO_2$ -eq/kg  $CH_4$ ) and to the higher fraction of concentrate in the diet (on average 1.26 instead 0.66 kg per L of milk) considered in Escribano *et al.* (2020), resulting finally in lower overall enteric  $CH_4$  emissions per ewe (8.64 versus 9.01 kg  $CH_4$  ewe<sup>-1</sup> year<sup>-1</sup>). On the other hand, when the emission intensity was referred to 1 ha of UAA and soil  $C_{seq}$  was not included, our results were in line with those of Escribano *et al.* (2020), where the emission intensity values were inversely related to the extent of permanent grasslands.

The percentage reduction of emission intensity with inclusion of soil  $C_{seq}$  observed in our study was consistent with results of Knudsen *et al.* (2019), who observed similar trends in dairy systems located in Western Europe. The inclusion of soil  $C_{seq}$  in the emission intensity estimate allowed to highlight better environmental performance in SE, regardless of the FU used. Indeed, emission intensity per kg FPCM and per ha of UAA in SE was 7% and 43% lower than SI, respectively, in accordance to what stated by other studies on extensive dairy sheep and goat systems based on permanent grasslands (Batalla *et al.*, 2015; O'Brien *et al.*, 2016; Gutiérrez-Peña *et al.*, 2019). Furthermore, Batalla *et al.* (2015) observed a significantly higher environmental performance of Mediterranean semi-intensive dairy sheep systems only when soil  $C_{seq}$  was excluded from the assessment. In their study, the inclusion of soil  $C_{seq}$ , estimated according to Petersen *et al.* (2013), determined an average decrease of 38% in emission intensity per kg of FPCM in sheep farms with Latxa breed. These farming systems were comparable with our case study in terms of stocking rate and feed supply. By contrast, in Escribano *et al.* (2020), extensive farming systems always showed the highest emission intensity per kg FPCM, even when soil  $C_{seq}$  was included. These contrasting results between Escribano *et al.* (2020) and our study may be due to the difference of soil  $C_{seq}$  values. In their study, in fact, the positive effect of soil  $C_{seq}$  from natural grasslands did not seem sufficient to compensate for the increase in enteric emissions of  $CH_4$ , due to the lower digestibility of the grass-based diet. On the other hand, in our study, the use of irrigated meadows and semi-natural grasslands, mainly based on legume species with high digestibility, certainly limited the negative effects of enteric  $CH_4$  emissions.

In our study, enteric  $CH_4$  emissions have been shown to be the largest contributor to GHG emissions, as demonstrated by several studies on ruminant livestock sector worldwide (González-García *et al.*, 2013; Marino *et al.*, 2016; O'Brien *et al.*, 2016). The different feeding strategies applied in the two production systems of our study influenced the environmental performance in terms of enteric emissions of  $CH_4$  per kg of FPCM. Indeed, despite the lower amount of fibrous feed ingested by ewes, the lower milk yield in SE (determined by lower DMI) resulted in higher enteric  $CH_4$  emissions per unit of FPCM than SI, with 67 versus 58 g of  $CH_4$  kg FPCM<sup>-1</sup>, respectively. As a consequence, enteric  $CH_4$  emissions had a larger contribution to the total emission intensity in SE than in SI (Table 7). In other terms, the FE of productive ewes in conjunction with digestibility represents the most important driver for enteric fermentation (Cottle *et al.*, 2011). Indeed, the value of FE was 15% higher in SI than in SE. The different contribution of pur-



**Figure 3.** Land use and soil C sequestration. Surface actually used to feed sheep (ha) and soil C sequestration (Mg  $CO_2$  per total surface occupied) deriving from crop residues, based on grassland type and use destination of the biomass, in the semi-intensive (A) and semi-extensive (B) system. The label values indicate the C sequestered (Mg) per whole surface of each grassland.

chased feeds and on-farm feeds indicated that feed supply chain and land use strategies strongly influenced the environmental performances of the two production systems, as already stated by Gislón *et al.* (2020).

In summary, the switch from SI to SE system resulted in a clear environmental advantage when emission intensity was referred to 1 ha of UAA, while the use of mass-based FU determined less pronounced differences between the two production systems (Salvador *et al.*, 2017; Escribano *et al.*, 2020). In this manner, we assessed the effect of the extensification with two contrasting perspectives and we observed that the use of less inputs per ha of UAA led to a clear environmental benefit. Considering that UAA was similar for both farming systems, this conclusion was not obvious and confirmed that using only a mass-based FU the environmental assessment did not provide a balanced view of the intensification impacts (Escribano *et al.*, 2020).

### Soil carbon sequestration

The values of soil  $C_{seq}$  per kg FPCM observed in our study were in line with the results obtained by Gutiérrez-Peña *et al.* (2019) in three different Spanish dairy goat systems, where  $C_{seq}$  per kg FPCM ranged from 0.15 to 0.81 kg CO<sub>2</sub>, with the highest values observed in extensive farms. Similarly, Escribano *et al.* (2020) showed analogous trend of soil  $C_{seq}$  moving from more intensive to more extensive Spanish dairy sheep farms, with values ranging between 0.09 and 2.04 kg CO<sub>2</sub> sequestered per kg of FPCM. The higher amount of soil  $C_{seq}$  in SE can be in part associated with the contribution of grazed grasslands, that occupied 83% of the UAA in SE compared to 35% in SI (Tables 1 and 3). For instance, according to Stanley *et al.* (2018), the increase of grazing surface together with the improvement of grazing management in SE contributed to enhance the soil  $C_{seq}$  of production system. The permanent grasslands and irrigated meadows also contributed considerably to the increase of soil  $C_{seq}$  in SE (Table 3). Thanks to their abundant root systems, high root turnover and rhizodeposition, the high amount of biomass residues left on the soil by these grasslands may explain higher soil  $C_{seq}$  values in SE (Beniston *et al.*, 2014; Lorenz and Lal, 2018). The increase of 'persistent' grasslands, such as irrigated meadows and permanent grasslands that persist undisturbed in the soil for a long time, could lead to an improvement of soil C input and soil C stock, if compared with temporary grasslands such as annual crops (King and Blesh, 2018; Gislón *et al.*, 2020). In addition, the soil disturbance caused by tillage in annual croplands can favour the soil organic matter mineralization (Six *et al.*, 2004; Acar *et al.*, 2018). Hence, the use of 'persistent' grasslands allowed to reduce the soil tillage intensity in SE. In this sense, Paustian *et al.* (1997) reported that management practices that can improve the soil C stock may increase the soil C input or decrease soil organic matter decomposition rates. Considering that a suitable forage system provides benefits in terms of soil  $C_{seq}$ , it can be considered as a climate change mitigation strategy of the whole dairy production system (Gislón *et al.*, 2020).

As mentioned before, similar studies conducted by Batalla *et al.* (2015), Gutiérrez-Peña *et al.* (2019) and Escribano *et al.* (2020) estimated soil  $C_{seq}$  of semi-extensive and semi-intensive Mediterranean small ruminant dairy systems adopting the model of Petersen *et al.* (2013). When soil  $C_{seq}$  was referred to 1 ha of UAA, these authors obtained similar values between production systems with different intensification levels. Contrarily, we estimated a soil  $C_{seq}$  value in SE nearly 70% greater than in SI. The values observed in our study were in line with the range of values showed by Gutiérrez-Peña *et al.* (2019), but they were lower than the average

values observed by Batalla *et al.* (2015) and higher than those showed by both Eldesouky *et al.* (2018) and Escribano *et al.* (2020). The cropping system productivity and the equations used for crop residues estimation can explain these different values of soil  $C_{seq}$ . The estimation of above- and belowground residues based on grassland type and destination use allowed to highlight the differences between SE and SI in terms of soil  $C_{seq}$  per ha, therefore confirming the potential of soil  $C_{seq}$  capacity of livestock systems based on permanent grasslands (Salvador *et al.*, 2017). Thus, the results of our study lead to suggest the need of detailed information about soil  $C_{seq}$  from cropping systems characterized by temporary and permanent grasslands, in LCA studies on small ruminant dairy farms. Obviously, it is essential that the models and coefficients used for estimating grazed biomass and crop residues are carefully adopted to obtain reliable soil  $C_{seq}$  values.

### Conclusions

Accounting for soil  $C_{seq}$  in emission intensity estimation, the transition from semi-intensive to semi-extensive Mediterranean dairy sheep farming led to a better environmental performance of the production system. By not counting soil  $C_{seq}$  within the LCA system boundaries, the adoption of semi-extensive system showed a clear environmental benefit only when emission intensity was expressed per ha of UAA, whereas semi-intensive systems resulted less impacting when emission intensity was referred to the kg of normalized milk.

Enteric CH<sub>4</sub> emissions were confirmed by far as the main source of GHG emissions with direct implications in feed supply strategies.

The soil  $C_{seq}$  was favoured by the presence of large areas covered by permanent grasslands and destined to the grazing in the semi-extensive production system. The improvement of soil organic C stock associated to the permanent grasslands would contribute effectively to mitigate GHG emissions in Mediterranean dairy sheep farms, highlighting the positive role of ecosystem services provided by extensive farming systems.

However, as the estimation of soil  $C_{seq}$  in our study was influenced by the methods used to estimate grazed biomass and crop residues, further investigations based on direct field measurements are advisable in order to improve data quality and results reliability.

### References

- Acar M, Celik I, Günel H, 2018. Effects of long-term tillage systems on aggregate-associated organic carbon in the eastern Mediterranean region of Turkey. *Eurasian J. Soil Sci.* 7:51-8.
- Agri-footprint 4.0, 2017. LCA database Blonk Agri-footprint, Gouda, Netherlands.
- Arzoumanidis I, Fullana-i-Palmer P, Raggi A, Gazulla C, Raugié M, Benveniste G, Anglada M, 2014. Unresolved issues in the accounting of biogenic carbon exchanges in the wine sector. *J. Clean. Prod.* 82:16-22.
- Atzori AS, Furesi R, Madau FA, Pulina P, Rattu PG, 2015. Sustainability of dairy sheep production in pasture lands: a case study approach to integrate economic and environmental perspectives. *Riv. Studi Sosten.* 1:117-34.
- Baldini C, Gardoni D, Guarino M, 2017. A critical review of the recent evolution of life cycle assessment applied to milk pro-

- duction. *J. Clean. Prod.* 140:421-35.
- Basset-Mens C, van der Werf HMG, 2005. Scenario-based environmental assessment of farming systems: the case of pig production in France. *Agric. Ecosyst. Environ.* 105:127-44.
- Batalla I, Knudsen MT, Mogensen L, del Hierro O, Pinto M, Hermansen JE, 2015. Carbon footprint of milk from sheep farming systems in Northern Spain including soil carbon sequestration in grasslands. *J. Clean. Prod.* 30:1-9.
- Beniston JW, DuPont ST, Glover JD, Lal R, Dungait JAJ, 2014. Soil organic carbon dynamics 75 years after land use change in perennial grassland and annual wheat agricultural systems. *Biogeochemistry* 120:37-49.
- Biddau R, Cidu R, 2005. Hydrogeochemical baseline studies prior to gold mining: a case study in Sardinia (Italy). *J. Geochem. Explor.* 86:61-85.
- Bolinder MA, Angers DA, Dubuc JP, 1997. Estimating shoot to root ratios and annual carbon inputs in soils for cereal crops. *Agric. Ecosyst. Environ.* 63:61-6.
- Bolinder MA, Janzen HH, Gregorich EG, Angers DA, VandenBygaart AJ, 2007. An approach for estimating net primary productivity and annual carbon inputs to soil for common agricultural crops in Canada. *Agric. Ecosyst. Environ.* 118:29-42.
- Brandão M, i Canals LM, 2013. Global characterisation factors to assess land use impacts on biotic production. *Int. J. Life Cycle Assess.* 18:1243-52.
- Burle M, Mielniczuk J, Focchi S, 1997. Effect of cropping systems on soil chemical characteristics, with emphasis on soil acidification. *Plant Soil.* 190:309-16.
- Chessa AP, Delitala A, 1997. Il clima della Sardegna. In: Collana Note Tecniche di Agrometeorologia per la Sardegna. Servizio agrometeorologico regionale per la Sardegna. Chiarella Ed., Sassari, Italy, pp 17-38.
- Cottle DJ, Nolan JV, Wiedemann SG, 2011. Ruminant enteric methane mitigation: a review. *Anim. Prod. Sci.* 51:491-514.
- Decandia M, Atzori AS, Acciaro M, Cabiddu A, Giovanetti V, Molina Alcaide E, Carro MD, Ranilla MJ, Molle G, Cannas A, 2011. Nutritional and animal factors affecting nitrogen excretion in sheep and goats. In: Ranilla MJ, Carro MD, Ben Salem H, Moran d-Fehr P (Eds.), *Challenging strategies to promote the sheep and goat sector in the current global context. Options Méditerranéennes: Série A. Séminaires Méditerranéens*; 99. CIHEAM/CSIC/Universidad de León/FAO, Zaragoza, Spain, pp 201-209.
- dos Santos NZ, Dieckow J, Bayer C, Molin R, Favaretto N, Pauletti V, Piva JT, 2011. Forages, cover crops and related shoot and root additions in no-till rotations to C sequestration in a subtropical Ferralsol. *Soil Tillage Res.* 111:208-18.
- EDA (European Dairy Association), 2018. Product environmental footprint category rules for dairy products. Available from: [http://ec.europa.eu/environment/eussd/smgp/pdf/PEFCR-DairyProducts\\_2018-04-25\\_V1.pdf](http://ec.europa.eu/environment/eussd/smgp/pdf/PEFCR-DairyProducts_2018-04-25_V1.pdf)
- Eldesouky A, Mesias FJ, Elghannam A, Escribano M, 2018. Can intensification compensate livestock greenhouse gas emissions? A study of the carbon footprint in Spanish agroforestry systems. *J. Clean. Prod.* 200:28-38.
- Escribano M, Elghannam A, Mesias FJ, 2020. Dairy sheep farms in semi-arid rangelands: a carbon footprint dilemma between intensification and land-based grazing. *Land Use Policy* 95:104600.
- Escudero A, González-Arias A, del Hierro O, Pinto M, Gartzia-Bengoetxea N, 2012. Nitrogen dynamics in soil amended with manures composted in dynamic and static systems. *J. Environ. Manage.* 108:66-72.
- FAO (Food and Agriculture Organization of the United Nations), 2010. Greenhouse gas emissions from the dairy sector - a life cycle assessment. Available from: <http://www.fao.org/docrep/012/k7930e/k7930e00.pdf>
- FAO (Food and Agriculture Organization of the United Nations), 2019. World food and agriculture - Statistical pocketbook. Available from: <http://www.fao.org/3/ca1796en/CA1796.EN.pdf>.
- Fedele A, Mazzi A, Niero M, Zuliani F, Scipioni A, 2014. Can the life cycle assessment methodology be adopted to support a single farm on its environmental impacts forecast evaluation between conventional and organic production? An Italian case study. *J. Clean. Prod.* 69:49-59.
- Flyvbjerg B, 2006. Five misunderstandings about case-study research. *Qual. Inq.* 12:219-45.
- Gerber PJ, Steinfeld H, Henderson B, Mottet A, Opio C, Dijkman J, Falcucci A, Tempio G, 2013. Tackling climate change through livestock: a global assessment of emissions and mitigation opportunities. Food and Agriculture Organization of the United Nations (FAO), Rome, Italy.
- Gislon G, Ferrero F, Bava L, Borreani G, Dal Prà A, Pacchioli MT, Sandrucci A, Zucali M, Tabacco E, 2020. Forage systems and sustainability of milk production: feed efficiency, environmental impacts and soil carbon stocks. *J. Clean. Prod.* 260:121012.
- González-García S, Hospido A, Moreira MT, Feijoo G, Arroja L, 2013. Environmental life cycle assessment of a Galician cheese: San Simon da Costa. *J. Clean. Prod.* 52:253-62.
- Gutiérrez-Peña R, Delgado-Pertiñez M, Manca C, Addis M, Guzmán-Guerrero JL, Sitzia M, 2018. Comparing lamb's meat quality of autochthonous sheep breeds Sarda and Mallorquina under different feeding management systems. In: 27<sup>th</sup> EGF General Meeting on 'Sustainable meat and milk production from grasslands. Cork, Ireland, 23:748-750.
- Gutiérrez-Peña R, Mena Y, Batalla I, Mancilla-Leytón JM, 2019. Carbon footprint of dairy goat production systems: A comparison of three contrasting grazing levels in the Sierra de Grazalema Natural Park (Southern Spain). *J. Environ. Manage.* 232:993-8.
- Hayashi K, Gaillard G, Nemecek T, 2006. Life cycle assessment of agricultural production systems: current issues and future perspectives. pp 98-109 in Proc of the International Seminar on Technology Development for Good Agriculture Practice (GAP) in Asia and Oceania. Food and Fertilizer Technology Center, Taipei, Taiwan, Japan.
- Horrillo A, Gaspar P, Díaz-Caro C, Escribano M, 2021. A scenario-based analysis of the effect of carbon pricing on organic livestock farm performance: a case study of Spanish dehesas and rangelands. *Sci. Total Environ.* 751:141675.
- IPCC (Intergovernmental Panel on Climate Change), 2007. The Physical Science Basis: Summary for Policymakers - Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Available from: <http://www.ipcc.ch/pdf/assessment-report/ar4/wg1/ar4-wg1-spm.pdf>
- IPCC (Intergovernmental Panel on Climate Change), 2013. Climate Change 2013: The Physical Science Basis - Working Group I Contribution to the Fifth Assessment Report of IPCC. Available from: <https://www.ipcc.ch/report/ar5/wg1/>
- IPCC (Intergovernmental Panel on Climate Change), 2019. Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories - Volume 4: Agriculture, forestry and other land use. Available from: <https://www.ipcc-nggip.iges.or.jp/public/2019rf/vol4.html>
- ISO (International Organization for Standardization), 2006a.

- 14040 - Environmental management - Life cycle assessment - Principles and framework. Standard number 14040:2006. ISO, Geneva, Switzerland.
- ISO (International Organization for Standardization), 2006b. 14044 - Environmental management - Life cycle assessment - Requirements and guidelines. Standard number 14040:2006. ISO, Geneva, Switzerland.
- ISPRA (Istituto Superiore per la Protezione e la Ricerca Ambientale); De Lauretis R, Romano D, Vitullo M, Arcarese C (Eds.), 2011. National greenhouse gas inventory system in Italy; Year 2011. Institute for Environmental Protection and Research, Environment Department, Rome, Italy.
- King AE, Blesh J, 2018. Crop rotations for increased soil carbon: perennality as a guiding principle. *Ecol. Appl.* 28:249-61.
- Knudsen MT, Dorca-Preda T, Djomo SN, Peña N, Padel S, Smith LG, Zollitsch W, Hörtenhuber S, Hermansen JE, 2019. The importance of including soil carbon changes, ecotoxicity and biodiversity impacts in environmental life cycle assessments of organic and conventional milk in Western Europe. *J. Clean. Prod.* 215:433-43.
- Lai R, Arca P, Lagomarsino A, Cappai C, Seddaiu G, Demurtas CE, Roggero PP, 2017. Manure fertilization increases soil respiration and creates a negative carbon budget in a Mediterranean maize (*Zea mays* L.)-based cropping system. *Catena* 151:202-12.
- Levasseur A, Lesage P, Margni M, Samson R, 2013. Biogenic carbon and temporary storage addressed with dynamic life cycle assessment. *J. Ind. Ecol.* 17:117-28.
- Lorenz K, Lal R, 2018. Carbon sequestration in grassland soils. In: *Carbon sequestration in agricultural ecosystems*. Springer, Cham., pp 175-209.
- Macciotta NPP, Cappio-Borlino A, Pulina G, 1999. Analysis of environmental effects on test day milk yields of Sarda dairy ewes. *J. Dairy Sci.* 82:2212-7.
- Marino R, Atzori AS, D'Andrea M, Iovane G, Trabalza-Marinucci M, Rinaldi L, 2016. Climate change: Production performance, health issues, greenhouse gas emissions and mitigation strategies in sheep and goat farming. *Small Rumin. Res.* 135:50-9.
- Mokany K, Raison RJ, Prokushkin AS, 2005. Critical analysis of root: shoot ratios in terrestrial biomes. *Glob. Change Biol.* 12:84-96.
- Moreno Ruiz E, Valsasina L, Fitzgerald D, Brunner F, Symeonidis A, Bourgault G, Wernet G, 2019. Documentation of changes implemented in the Ecoinvent database v3.6. Ecoinvent Association, Zürich, Switzerland. Available from: [https://www.ecoinvent.org/files/change\\_report\\_v3\\_6\\_20190912.pdf](https://www.ecoinvent.org/files/change_report_v3_6_20190912.pdf)
- Nayak AK, Rahman MM, Naidu R, Dhal B, Swain CK, Nayak AD, Tripathi R, Shahid M, Islam MR, Pathak H, 2019. Current and emerging methodologies for estimating carbon sequestration in agricultural soils: a review. *Sci. Total Environ.* 665:890-912.
- Nemecek T, Kägi T, 2007. Life Cycle Inventories of Swiss and European Agricultural Production Systems. Final report ecoinvent V2.0 No. 15a. Agroscope Reckenholz-Taenikon Research Station ART, Swiss Centre for Life Cycle Inventories, Zurich and Dübendorf, Switzerland. Available from: [www.ecoinvent.ch](http://www.ecoinvent.ch)
- Notarnicola B, Sala S, Anton A, McLaren SJ, 2017. The role of life cycle assessment in supporting sustainable agri-food systems: A review of the challenges. *J. Clean. Prod.* 140:399-409.
- O'Brien D, Bohan A, McHugh N, Shaloo L, 2016. A life cycle assessment of the effect of intensification on the environmental impacts and resource use of grass-based sheep farming. *Agric. Syst.* 148:95-104.
- Paustian K, Andren O, Janzen HH, Lal R, Smith P, Tian G, Tiessen H, Van Noordwijk M, Woomer PL, 1997. Agricultural soils as a sink to mitigate CO<sub>2</sub> emissions. *Soil Use Manag.* 13:230-44.
- Petersen BM, 2010. A model for the carbon dynamics in agricultural, mineral soils. Technical Report. Aarhus University, Denmark. Available from: [www.agrsci.dk/c-tool](http://www.agrsci.dk/c-tool)
- Petersen BM, Knudsen MT, Hermansen JE, Halberg N, 2013. An approach to include soil carbon changes in life cycle assessments. *J. Clean. Prod.* 52:217-24.
- Pirlo G, Carè S, Fantin V, Falconi F, Buttol P, Terzano GM, Masoni P, Pacelli P, 2014. Factors affecting life cycle assessment of milk produced in 6 Mediterranean buffalo farms. *J. Dairy Sci.* 97:6583-93.
- Porqueddu C, Melis RAM, Franca A, Sanna F, Hadjigeorgiou I, Casasús I, 2017. The role of grasslands in the less favoured areas of Mediterranean Europe. In: 19<sup>th</sup> EGF Symposium on 'Grassland resources for extensive farming systems in marginal lands: major drivers and future scenarios - Grassland Science in Europe'. Alghero, Sardinia, Italy, 22:3-22.
- PRé Consultants, 2018. Software LCA SimaPro Analyst 8.5.0.0.
- Pulina G, Nudda A, 2002. Milk production. In: *Dairy sheep feeding and nutrition*. Avenue media (ed.), Bologna, Italy, pp 11-13.
- Pulina G, Milán MJ, Lavín MP, Theodoridis A, Morin E, Capote J, Thomas DL, Francesconi AHD, Caja G, 2018. Invited review: current production trends, farm structures, and economics of the dairy sheep and goat sectors. *J. Dairy Sci.* 101:6715-29.
- Rural Development Programme of Sardinia, 2014-2020. Programma di sviluppo rurale PSR Sardegna. Available from: <http://www.regione.sardegna.it/speciali/programmasvilupporurale/benvenuto-sul-sito-del-psr-2014-2020>
- Salou T, Le Mouël C, van der Werf HMG, 2017. Environmental impacts of dairy system intensification: the functional unit matters! *J. Clean. Prod.* 140:445-54.
- Salvador S, Corazzin M, Romanzin A, Bovolenta S, 2017. Greenhouse gas balance of mountain dairy farms as affected by grassland carbon sequestration. *J. Environ. Manage.* 196:644-50.
- Sardegna Agricoltura, 2013. Note tecniche sull'alimentazione degli ovini e dei caprini: Opuscolo n. 2. Available from: [http://www.sardegnaagricoltura.it/documenti/14\\_43\\_20131002143844.pdf](http://www.sardegnaagricoltura.it/documenti/14_43_20131002143844.pdf)
- Searates, 2021. Shipping to and from anywhere in the world. Find the best freight quote. Available from: <https://www.searates.com>
- Seddaiu G, Bagella S, Pulina A, Cappai C, Salis L, Rossetti I, Lai R, Roggero PP, 2018. Mediterranean cork oak wooded grasslands: synergies and trade-offs between plant diversity, pasture production and soil carbon. *Agrofor. Syst.* 92:893-908.
- Six J, Bossuyt H, Degryzed S, Denef K, 2004. A history of research on the link between (micro)aggregates, soil biota, and soil organic matter dynamics. *Soil Tillage Res.* 79:7-31.
- Stanley PL, Rowntree JE, Beede DK, DeLonge MS, Hamm MV, 2018. Impacts of soil carbon sequestration on life cycle greenhouse gas emissions in Midwestern USA beef finishing systems. *Agric. Syst.* 162:249-58.
- Tedeschi LO, Cannas A, Fox DG, 2008. A nutrition mathematical model to account for dietary supply and requirements of energy and nutrients for domesticated small ruminants: the development and evaluation of the small ruminant nutrition system. *R. Bras. Zootec.* 37:178-90.

Vagnoni E, Franca A, Porqueddu C, Duce P, 2017. Environmental profile of Sardinian sheep milk cheese supply chain: A comparison between two contrasting dairy systems. *J. Clean. Prod.* 165:1078-89.

Vermorel M, Jouany JP, Eugène M, Sauvant D, Noblet J, Dourmad JY, 2008. Evaluation quantitative des émissions de méthane entérique par les animaux d'élevage en 2007 en France. *INRA Prod. Anim.* 21:403-18.

Non-commercial use only